





What was your last PSA blood test for prostate cancer? \_\_\_\_\_ When was it drawn? \_\_\_\_\_

Please circle only one response for each answer	Not at all	Less than 1 time in 5	Less than half the time	About half the time	More than half the time	Almost always
1. Over the past month or so, how often have you had the sensation of not emptying your bladder completely after you finish urinating?	0	1	2	3	4	5
2. Over the past month or so, how often have you had to urinate again less than two hours after you finished urinating?	0	1	2	3	4	5
3. Over the past month or so, how often have you stopped and started again several times when you urinated?	0	1	2	3	4	5
4. Over the past month or so, how often have you found it difficult to postpone urination?	0	1	2	3	4	5
5. Over the past month or so, how often have you had a weak urinary stream?	0	1	2	3	4	5
6. Over the past month or so, how often have you had to push or strain to begin urination?	0	1	2	3	4	5
7. Over the past month or so, how many times did you most typically get up to urinate from the time you went to bed until the time you got up in the morning?	0	1	2	3	4	5

If you were to spend the rest of your life with your urinary condition just as it is now, how would you feel about that?

Delighted    Pleased    Mostly satisfied    Mixed    Mostly dissatisfied    Unhappy    Terrible

If you have problems with your erections, please answer the following questions; if not, go to the last page.

1. How do you rate your **confidence** that you could **keep** an erection?

Very low 1	Low 2	Moderate 3	High 4	Very high 5
---------------	----------	---------------	-----------	----------------

2. When you had erections with sexual stimulation, **how often** were your erections hard enough for penetration (entering your partner)?

No sexual activity 0	Almost never or never 1	A few times (much less than half the time) 2	Sometimes (about half the time) 3	Most times (much more than half) 4	Almost always or always 5
-------------------------	----------------------------	---	--------------------------------------	---------------------------------------	------------------------------

3. During sexual intercourse **how often** were you able to maintain your erection after you had penetrated (entered) your partner?

Did not attempt intercourse 0	Almost never or never 1	A few times (much less than half the time) 2	Sometimes (about half the time) 3	Most times (much more than half) 4	Almost always or always 5
----------------------------------	----------------------------	---	--------------------------------------	---------------------------------------	------------------------------

4. During sexual intercourse, **how difficult** was it to maintain your erection to completion?

Did not attempt intercourse 0	Extremely difficult 1	Very difficult 2	Difficult 3	Slightly difficult 4	Not difficult 5
----------------------------------	--------------------------	---------------------	----------------	-------------------------	--------------------

5. When you attempted sexual intercourse, **how often** was it satisfactory for you?

Did not attempt intercourse 0	Almost never or never 1	A few times (much less than half the time) 2	Sometimes (about half the time) 3	Most times (much more than half) 4	Almost always or always 5
----------------------------------	----------------------------	---	--------------------------------------	---------------------------------------	------------------------------

## Review of Systems

Please check off any of the items below that may have been a problem for you recently.

- Constitutional**     chronic fever     joint pain     weight loss > 20 lb.  
 night sweats     bone pain     recent severe pain
- Eyes**     glaucoma     blurry vision     cataracts     double vision     retinitis pigmentosa
- Ear/Nose/Throat**     earaches     dizziness     chronic sore throat  
 ringing in ears     nose bleeds     hoarseness
- Cardiovascular**     heart attack     heart failure     heart murmur     irregular heart beat  
 atrial fibrillation     palpitations     rheumatic fever     artificial heart valve  
 chest pain     chest pressure     chest tightness     pain in legs when walking  
 leg swelling     use nitroglycerin on exertion
- Respiration**     shortness of breath     pneumonia     emphysema     chronic cough  
 coughing up blood     asthma     wheezing     tuberculosis
- Gastrointestinal**     appetite     hepatitis     bloody stools     ulcers  
 nausea     liver trouble     diarrhea     heartburn  
 vomiting     jaundice     incontinence of stool     constipation  
 gallstones     hiatal hernia
- Genitourinary**     venereal disease     split urinary stream     kidney stones     incontinence  
 sexually transmitted disease     protein in urine     sugar in urine     trouble with ejaculation  
 urinary strictures     blood in urine     sex drive     kidney infection  
 infertility     epididymitis     prostatitis
- Musculoskeletal**     arthritis     back pain     disc disease  
 muscle weakness     artificial joints
- Skin**     unexplained rashes     redness     hives     moles
- Neurological**     seizures     numbness     stroke     difficulty walking  
 tingling     memory loss     mini-stroke     tremor
- Psychiatric**     schizophrenia     anxiety     paranoia     depression     bipolar disorder
- Endocrine**     diabetes     weight gain     increased thirst     growth delays  
 thyroid trouble     weight loss     breast lumps     late puberty
- Hematologic/Lymphatic/Immunologic**     swollen lymph nodes     easy bruising     sickle cell  
 lupus     anemia     nosebleeds  
 abnormal bleeding     mumps     HIV/AIDS